

**Math 800 Final Exam**  
**Spring 2017**

**Note:** These 10 problems are taken from the complex analysis part of UCLA qualifying exams in Analysis for fall 2016 and spring 2016 at <https://secure.math.ucla.edu/gradquals/hbquals.php>. Due by Wednesday, May 3rd in class.

1) Determine

$$\int_0^\infty \frac{x^{a-1}}{x+z} dx$$

for  $0 < a < 1$  and  $\operatorname{Re} z > 0$ . Justify all manipulations.

2) Let  $\mathbf{C}_+ = \{z \in \mathbf{C} : \operatorname{Im} z > 0\}$  and let  $f_n : \mathbf{C}_+ \rightarrow \mathbf{C}_+$  be a sequence of holomorphic functions. Show that unless  $|f_n| \rightarrow \infty$  uniformly on compact subsets of  $\mathbf{C}_+$ , there exists a subsequence converging uniformly on compact subsets of  $\mathbf{C}_+$ .

3) Let  $f : \mathbf{C} \rightarrow \mathbf{C}$  be entire and assume that  $|f(z)| = 1$  when  $|z| = 1$ . Show that  $f$  is in the following form,

$$f(z) = Cz^m$$

for some integer  $m > 0$  and  $C \in \mathbf{C}$  with  $|C| = 1$ .

4) Does there exist a function  $f(z)$  holomorphic in the disk  $|z| < 1$  such that  $\lim_{|z| \rightarrow 1} |f(z)| = \infty$ ? Either find one or prove that none exist.

5) Assume that  $f(z)$  is holomorphic on  $|z| < 2$ . Show that

$$\max_{|z|=1} f(z) - \frac{1}{z} \geq 1.$$

6)

(a). Find a real-valued harmonic function  $v$  defined on the disk  $|z| < 1$  such that  $v(z) > 0$  and  $\lim_{z \rightarrow 1} v(z) = \infty$ .

(b). Let  $u$  be a real-valued harmonic function in the disk  $|z| < 1$  such that  $u(z) \leq M < \infty$  and  $\limsup_{r \rightarrow 1} u(re^{i\theta}) \leq 0$  for all  $\theta \in (0, 2\pi)$ . Show that  $u(z) \leq 0$ . The function in part (a) is useful here.

7) Let  $\mathcal{H}$  be the space of holomorphic functions  $f$  in  $D = \{z \in \mathbf{C} : |z| < 1\}$  such that

$$\int_D |f(z)|^2 dA(z) < \infty,$$

where the integration is with respect to the Lebesgue measure  $A$  on  $D$ . The vector space  $\mathcal{H}$  is a Hilbert space if equipped with the inner product

$$\langle f, g \rangle = \int_D f(z) \bar{g}(z) dA(z)$$

for  $f, g \in \mathcal{H}$ . Fix  $z_0 \in D$  and define  $L_{z_0}(f) = f(z_0)$  for  $f \in \mathcal{H}$ .

(a). Show that  $L_{z_0}$  is a bounded linear functional on  $\mathcal{H}$ .

(b). Find an explicit  $g_{z_0} \in \mathcal{H}$  such that

$$L_{z_0}(f) = \langle f, g_{z_0} \rangle$$

for all  $f \in \mathcal{H}$ .

**8)** Let  $f$  be a continuous complex-valued function on the closed unit disk  $\bar{D} = \{z \in \mathbf{C} : |z| \leq 1\}$  such that  $f$  is holomorphic in the open disk  $D = \{z \in \mathbf{C} : |z| < 1\}$  and  $f(0) \neq 0$ . Show that

(a). Let  $0 < r < 1$  and  $\inf_{|z|=r} |f(z)| > 0$ . Then

$$\frac{1}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta \geq \log |f(0)|.$$

(b). The measure  $|\{\theta \in [0, 2\pi] : f(e^{i\theta}) = 0\}| = 0$ , where  $|E|$  denotes the Lebesgue measure of  $E \subset [0, 2\pi]$ .

**9)** Let  $\mu$  be a positive Borel measure on  $[0, 1]$  with  $\mu([0, 1]) = 1$ .

(a). Show that the function  $f(z)$  defined as

$$f(z) = \int_{[0,1]} e^{izt} d\mu(t)$$

for  $z \in \mathbf{C}$  is holomorphic on  $\mathbf{C}$ .

(b). Suppose that there exists  $n \in \mathbf{N}$  such that

$$\limsup_{|z| \rightarrow \infty} \frac{|f(z)|}{|z|^n} < \infty.$$

Show that  $\mu$  equals the Dirac measure  $\delta_0$  at 0.

**9)** Show that  $f : \mathbf{C} \rightarrow \mathbf{C}$  is holomorphic function such that the function

$$z \mapsto g(z) = f(z)f(1/z)$$

is bounded on  $\mathbf{C} \setminus \{0\}$ .

(a). Show that if  $f(0) \neq 0$ , then  $f$  is a constant.

(b). Show that if  $f(0) = 0$ , then there exists  $a \in \mathbf{C}$  and  $n \in \mathbf{N}$  such that  $f(z) = az^n$  for all  $z \in \mathbf{C}$ .